



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

journal homepage: www.elsevier.com/locate/jmaa



Regular Articles

Viscosity-flux approximation method to inhomogeneous system of isentropic gas dynamics [☆]



Xianting Wang ^a, Yun-Guang Lu ^{b,*}, Christian Klingenberg ^c

^a *Wuxi Institute of Technology, China*

^b *School of Mathematics, Hangzhou Normal University, China*

^c *Department of Mathematics, Wuerzburg University, Germany*

ARTICLE INFO

Article history:

Received 1 February 2023
Available online 20 October 2023
Submitted by D. Donatelli

Keywords:

Global L^∞ solution
Inhomogeneous system of isentropic gas dynamics
Flux approximation
Compensated compactness

ABSTRACT

In this paper, we apply the maximum principle with the viscosity-flux approximation to obtain the a-priori L^∞ estimates on the Riemann invariants of (1.1), $\Delta(\nu^{\xi,\varepsilon}, u^{\xi,\varepsilon}) \leq \omega(x - kt)$ and $\Upsilon(\nu^{\xi,\varepsilon}, u^{\xi,\varepsilon}) \leq \omega(x - kt)$ for the solutions $(\nu^{\xi,\varepsilon}, u^{\xi,\varepsilon})$ of the Cauchy problem (1.6) and (1.7), where $\omega(x - kt)$ is a nonnegative bounded function, and to prove the global existence of the L^∞ entropy solutions for the Cauchy problem of inhomogeneous system of isentropic gas dynamics (1.1) with arbitrary bounded initial data (1.2).

© 2023 Elsevier Inc. All rights reserved.

1. Introduction

In this paper we studied the Cauchy problem of the following system of isentropic gas dynamics with a moving source term

$$\begin{cases} \nu_t + (\nu u)_x = \alpha(x - kt)\nu u \\ (\nu u)_t + (\nu u^2 + \sigma(\nu))_x = \alpha(x - kt)\nu u^2 - \zeta(x, t)\nu u|u|, \end{cases} \quad (1.1)$$

where u is the velocity of gas, ν the density, $\sigma(\nu)$ the pressure, the positive constant k denotes a moving speed [8], and $\zeta(x, t) \geq 0$ is a friction function of the space variable x and the time variable t . In physics, the pressure σ taking the special form $\sigma(\nu) = \frac{1}{\gamma}\nu^\gamma$, is for the polytropic gas, where the adiabatic exponent $\gamma > 1$.

[☆] The authors thank the anonymous reviewers for their valuable comments. This paper is supported by the NSFC (12071106) of China and a Humboldt renewed research fellowship of Germany.

* Corresponding author.

E-mail addresses: wangxt@wxit.edu.cn (X.T. Wang), yglu2010@hznu.edu.cn (Y.-G. Lu), klingen@mathematik.uni-wuerzburg.de (C. Klingenberg).

When ζ is a constant, $k = 0$ and $\alpha(x) = -\frac{a'(x)}{a(x)}$, where the function $a(x)$ denotes a slowly variable cross section area at x in the nozzle, the global entropy solutions for the Cauchy problem (1.1) with bounded initial data

$$(\nu(x, 0), u(x, 0)) = (\nu_0(x), u_0(x)), \quad \nu_0(x) \geq 0 \tag{1.2}$$

were first studied in [1,14] when the adiabatic exponent $1 < \gamma \leq \frac{5}{3}$, and by the author in [11] for any adiabatic exponent $\gamma > 1$, if the initial data satisfy the strong restriction condition $\Upsilon(\nu_0(x), u_0(x)) \leq 0$, where Υ is a Riemann invariant given in (1.8). The initial-boundary value problem of compressible Euler equations including friction and heating that model the transonic Fanno-Rayleigh flows through symmetric variable area nozzles is studied in [2].

It is well-known that the unique difficulty to deal with the inhomogeneous system (1.1) is to obtain the a-priori L^∞ estimates of the approximation solutions of (1.1), for instance, the a-priori L^∞ estimates of the classical viscosity solutions for the Cauchy problem of the following parabolic system

$$\begin{cases} \nu_t + (\nu u)_x = \alpha(x - kt)\nu u + \varepsilon\nu_{xx} \\ (\nu u)_t + (\nu u^2 + \sigma(\nu))_x = \alpha(x - kt)\nu u^2 - \zeta(x, t)\nu u|u| + \varepsilon(\nu u)_{xx}, \end{cases} \tag{1.3}$$

with the initial data (1.2).

When $\alpha(x - kt) = 0$, (1.1) is the river flow equations, a shallow-water model describing the vertical depth ν and mean velocity u , where $\zeta(x, t)\nu u|u|$ corresponds physically to a friction term and ζ is the friction coefficient. This kind of inhomogeneous systems is simple since the source terms have, in some senses, the symmetric behavior (see [4] for the details).

When $\zeta(x, t) = 0$ and $k = 0$, i.e., the nozzle flow without friction, system (1.1) was well studied in (cf. [1,9,10,12,15–17], and the references cited therein). Roughly speaking, the technique introduced in these papers, is to control the super-linear source terms $\alpha(x)\nu u$ and $\alpha(x)\nu u^2$ by the flux functions νu and $\nu u^2 + \sigma(\nu)$ in (1.1) and to deduce a upper bound of Δ or Υ by a bounded nonnegative function $\psi(x)$, which depends on the function $\alpha(x)$.

When $\alpha(x) \not\equiv 0$ and $\zeta(x, t) \not\equiv 0$, both the above techniques do not work because the flux functions can not be used to control the super-linear friction source terms $\zeta\nu u|u|$, and the functions $\alpha(x)\nu u$ destroy the symmetry of the Riemann invariants (Δ, Υ) (see [5] for the numerical analysis).

However, we may copy the following steps given in [12] to overcome the above difficulty.

First, to avoid the singularity of the flux function νu^2 near the vacuum $\nu = 0$, we still use the technique of the ξ -flux-approximation given in [13] and introduce the sequence of systems

$$\begin{cases} \nu_t + (-2\xi u + \nu u)_x = \alpha(x - kt)(\nu - 2\xi)u \\ (\nu u)_t + (\nu u^2 - \xi u^2 + \sigma_1(\nu, \xi))_x = \alpha(x - kt)(\nu - 2\xi)u^2 - \zeta(x, t)\nu u|u| \end{cases} \tag{1.4}$$

to approximate system (1.1), where $\xi > 0$ denotes a regular perturbation constant and the perturbation pressure

$$\sigma_1(\nu, \xi) = \int_{2\xi}^{\nu} \frac{t - 2\xi}{t} \sigma'(t) dt. \tag{1.5}$$

Second, we add the classical viscosity terms to the right-hand side of (1.4) and obtain the following standard parabolic system

$$\begin{cases} \nu_t + ((\nu - 2\xi)u)_x = \alpha(x - kt)(\nu - 2\xi)u + \varepsilon\nu_{xx} \\ (\nu u)_t + (\nu u^2 - \xi u^2 + \sigma_1(\nu, \xi))_x = \alpha(x - kt)(\nu - 2\xi)u^2 - \zeta(x, t)\nu u|u| + \varepsilon(\nu u)_{xx} \end{cases} \tag{1.6}$$

with initial data

$$(\nu^{\xi, \varepsilon}(x, 0), u^{\xi, \varepsilon}(x, 0)) = (\nu_0(x) + 2\xi, u_0(x)), \tag{1.7}$$

where $(\nu_0(x), u_0(x))$ are given in (1.2). Now multiplying (1.6) by $(\frac{\partial \Delta}{\partial \nu}, \frac{\partial \Delta}{\partial m})$ and $(\frac{\partial \Upsilon}{\partial \nu}, \frac{\partial \Upsilon}{\partial m})$ respectively, where

$$\Upsilon(\nu, u) = \int_c^\nu \frac{\sqrt{\sigma'(s)}}{s} ds - u, \quad \Delta(\nu, u) = \int_c^\nu \frac{\sqrt{\sigma'(s)}}{s} ds + u \tag{1.8}$$

are the Riemann invariants of (1.1), c is a constant and $m = \nu u$ denotes the momentum, we obtain

$$\begin{aligned} &\Delta_t + \Lambda_2^\xi \Delta_x \\ &= \varepsilon \Delta_{xx} + \frac{2\varepsilon}{\nu} \nu_x \Delta_x - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') \nu_x^2 \\ &+ \alpha(x - kt)(\nu - 2\xi)u \frac{\sqrt{\sigma'(\nu)}}{\nu} - \zeta(x, t)u|u| \end{aligned} \tag{1.9}$$

and

$$\begin{aligned} &\Upsilon_t + \Lambda_1^\xi \Upsilon_x \\ &= \varepsilon \Upsilon_{xx} + \frac{2\varepsilon}{\nu} \nu_x \Upsilon_x - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') \nu_x^2 \\ &+ \alpha(x - kt)(\nu - 2\xi)u \frac{\sqrt{\sigma'(\nu)}}{\nu} + \zeta(x, t)u|u|, \end{aligned} \tag{1.10}$$

where

$$\Lambda_1^\xi = \frac{m}{\nu} - \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}, \quad \Lambda_2^\xi = \frac{m}{\nu} + \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \tag{1.11}$$

are two eigenvalues of the approximation system (1.4).

It is obvious that the terms $\alpha(x - kt)(\nu - 2\xi)u \frac{\sqrt{\sigma'(\nu)}}{\nu}$ in (1.9) and (1.10) are not symmetric with respect to the Riemann invariants Δ, Υ . However, with the strong restriction $\Upsilon(\nu_0(x), u_0(x)) \leq 0$ on the initial data, we obtained the uniformly upper bounds of Υ and Δ in [14,11], by using the maximum principle. Unfortunately, without the condition $\Upsilon(\nu_0(x), u_0(x)) \leq 0$, we will meet some new technical difficulties when we study the Cauchy problem (1.1) and (1.2).

2. Main results

In this paper, we will unite the techniques given in [4] and in [14,11] to obtain the estimates $\Upsilon \leq \omega(x - kt)$ and $\Delta \leq \omega(x - kt)$, for a suitable uniformly bounded function $\omega(x - kt)$, and to prove the global existence of the entropy solutions for the Cauchy problem (1.1) and (1.2).

The main results of this paper are in the following Theorems 1-3.

Theorem 1. Let $\sigma(\nu) = \frac{1}{\gamma} \nu^\gamma, \gamma > 1, \zeta(x, t) \geq 0$ and $\alpha(x - kt) \leq 0$.

(I). Let $1 < \gamma \leq 3, k > 0$. Then there exists a function $\psi(x) \in \mathcal{B}_d^2(-\infty, \infty)$ satisfying $\psi(x) \leq M < k$ and the inequality (2.5) given below such that

$$\Upsilon(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) = \frac{(\nu^{\xi,\varepsilon}(x, t))^\theta}{\theta} - u^{\xi,\varepsilon}(x, t) \leq \psi(x - kt) \tag{2.1}$$

and

$$\Delta(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) = \frac{(\nu^{\xi,\varepsilon}(x, t))^\theta}{\theta} + u^{\xi,\varepsilon}(x, t) \leq \psi(x - kt) \tag{2.2}$$

if the initial data $\Upsilon(\nu^{\xi,\varepsilon}(x, 0), u^{\xi,\varepsilon}(x, 0)) \leq \psi(x)$ and $\Delta(\nu^{\xi,\varepsilon}(x, 0), u^{\xi,\varepsilon}(x, 0)) \leq \psi(x) \in \mathcal{B}_d^2$, where $\psi_x = \frac{\partial \psi(x-kt)}{\partial x}$ and $\theta = \frac{\gamma-1}{2}$.

(II). Let $\gamma > 3, k < 0$. Then there exists a function $\chi(x) \in \mathcal{C}_d^2(-\infty, \infty)$ satisfying $\chi(x) \leq M < -\frac{2}{\gamma+1}k$ and the inequality (2.6) given below such that $\Upsilon(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) \leq \chi(x - kt)$, $\Delta(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) \leq \chi(x - kt)$, which are similar with the estimates (2.1) and (2.2).

Theorem 2. Let $\sigma(\nu) = \frac{1}{\gamma}\nu^\gamma, \gamma > 1, \zeta(x, t) \geq 0$ and $\alpha(x - kt) = \alpha_-(x - kt) + \alpha_+(x - kt)$, where $\alpha_-(x - kt) \leq 0, \alpha_+(x - kt) \geq 0$.

(III). Let $1 < \gamma \leq 3, k > 0$. Then there exists a function $\psi(x) \in \mathcal{B}_d^2(-\infty, \infty)$ satisfying $\psi(x) \leq M < k$, the inequalities (2.12) and (2.13) given below, such that the same estimates like (2.1) and (2.2) are true.

(IV). Let $\gamma > 3, k < 0$. Then there exists a function $\chi(x) \in \mathcal{C}_d^2(-\infty, \infty)$ satisfying $\chi(x) \leq M < -\frac{2}{\gamma+1}k$, the inequalities (2.14) and (2.15) given below, such that $\Upsilon(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) \leq \chi(x - kt)$, $\Delta(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) \leq \chi(x - kt)$ are true.

Theorem 3. If the conditions about the functions $\alpha(x - kt), \zeta(x, t)$ and the initial data in Theorem 1 or Theorem 2 are satisfied, then there exists a subsequence of $(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t))$, which converges pointwisely to a pair of bounded functions $(\nu(x, t), u(x, t))$ as ξ, ε tend to zero, and the limit is a weak entropy solution of the Cauchy problem (1.1)-(1.2).

Definition 1. We call a pair of bounded functions $(\nu(x, t), u(x, t))$ is a weak entropy solution of the Cauchy problem (1.1)-(1.2) if

$$\begin{cases} \int_0^\infty \int_{-\infty}^\infty \nu \phi_t + (\nu u) \phi_x + \alpha(x - kt) \nu u \phi dx dt + \int_{-\infty}^\infty \nu_0(x) \phi(x, 0) dx = 0, \\ \int_0^\infty \int_{-\infty}^\infty \nu u \phi_t + (\nu u^2 + \sigma(\nu)) \phi_x + (\alpha(x - kt) \nu u^2 - \zeta(x, t) \nu u |u|) \phi dx dt \\ + \int_{-\infty}^\infty \nu_0(x) u_0(x) \phi(x, 0) dx = 0 \end{cases} \tag{2.3}$$

holds for all test function $\phi \in \mathcal{C}_0^1(R \times R^+)$ and

$$\begin{aligned} & \int_0^\infty \int_{-\infty}^\infty \eta(\nu, m) \phi_t + q(\nu, m) \phi_x + \alpha(x - kt) \nu u \eta(\nu, m)_\nu \\ & + (\alpha(x - kt) \nu u^2 - \zeta(x, t) \nu u |u|) \eta(\nu, m)_m \phi dx dt \geq 0 \end{aligned} \tag{2.4}$$

holds for any non-negative test function $\phi \in \mathcal{C}_0^\infty(R \times R^+ - \{t = 0\})$, where $m = \nu u$ and (η, q) is a pair of convex entropy-entropy flux of system (1.1).

Remark 1. The definitions of $\mathcal{B}_d^2(R)$ and $\mathcal{C}_d^2(R)$ are given in [11]. For a given $\psi(x) \in \mathcal{B}_d^2(R)$, if we let $\chi(x) = \psi(-x)$, then $\chi(x) \in \mathcal{C}_d^2(R)$.

Before we prove Theorems 1-3 in the next sections, we first construct some necessary special functions in $\mathcal{B}_d^2(R)$ and $\mathcal{C}_d^2(R)$.

Example 1. For a given $\alpha(x)$, we can find many functions $\psi(x)$ in the set $\mathcal{B}_d^2(R)$, which satisfy

$$\begin{aligned} &\theta^2\alpha^2(x-kt)\psi^2(x-kt) + (1-\theta)^2\psi_x^2 - 2\theta(1+\theta)\alpha(x-kt)\psi(x-kt)\psi_x \\ &+ 4\varepsilon_1\theta\alpha(x-kt)\psi(x-kt)\psi_x \leq 0 \end{aligned} \tag{2.5}$$

or $\chi(x)$ in $\mathcal{C}_d^2(R)$ satisfy

$$\begin{aligned} &\theta^2\alpha^2(x-kt)\chi^2(x-kt) + (1-\theta)^2\chi_x^2 + 2\theta(1+\theta)\alpha(x-kt)\chi(x-kt)\chi_x \\ &- 4\varepsilon_1\theta\alpha(x-kt)\chi(x-kt)\chi_x \leq 0, \end{aligned} \tag{2.6}$$

for a small $\varepsilon_1 > 0$.

For instance, we choose $a(x) = x^2, \alpha(x) = -\frac{a'(x)}{a(x)} = -\frac{2}{x}$ as the author studied in [15] for the spherically, symmetric solutions in $x > 1$. We now extend it to the whole space $x \in (-\infty, \infty)$ in the following way:

$$\alpha(x) = \begin{cases} -\frac{2}{x}, & \text{for } x > \varepsilon_0, \\ -\frac{2x}{\varepsilon_0^2}, & \text{for } 0 \leq x \leq \varepsilon_0, \\ 0, & \text{for } x < 0, \end{cases} \tag{2.7}$$

where $\varepsilon_0 > 0$ is a constant. Then we can easily check that the following function

$$\psi(x) = \begin{cases} qx^\beta, & \text{for } x > \varepsilon_0, \\ q_1e^{\frac{\beta x^2}{2\varepsilon_0^2}}, & \text{for } 0 \leq x \leq \varepsilon_0, \\ q_1, & \text{for } x < 0, \end{cases} \tag{2.8}$$

satisfies

$$\psi'(x) = -\frac{\beta}{2}\alpha(x)\psi(x) \text{ if } q\varepsilon_0^\beta = q_1e^{\frac{\beta}{2}}, \tag{2.9}$$

where q, q_1 are two positive constants and β is a negative constant. Then clearly $-M\psi(x) \leq \psi'(x) \leq 0$.

Moreover,

$$\psi''(x) = \begin{cases} q\beta(\beta-1)x^{\beta-2}, & \text{for } x > \varepsilon_0, \\ q_1\frac{\beta}{\varepsilon_0^2}e^{\frac{\beta x^2}{2\varepsilon_0^2}} + q_1\left(\frac{\beta x}{\varepsilon_0^2}\right)^2e^{\frac{\beta x^2}{2\varepsilon_0^2}}, & \text{for } 0 \leq x \leq \varepsilon_0, \\ 0, & \text{for } x < 0, \end{cases} \tag{2.10}$$

and $\psi''(x) = \psi_1(x) + \psi_2(x)$, where $\psi_1(x), \psi_2(x)$ satisfy the conditions in the definition of $\mathcal{B}_d^2(R)$, thus $\psi(x) \in \mathcal{B}_d^2(R)$.

Let $-\frac{\beta}{2} = \frac{\theta}{1-\theta}$. Then we can easily check that $\psi(x-kt)$ satisfies (2.5), where $\psi(x)$ is given by (2.9).

In fact, let $\psi_x = \frac{\partial\psi(x-kt)}{\partial x} = \frac{\theta}{1-\theta}\alpha(x-kt)\psi(x-kt)$. Then $\psi_x \leq 0$ and (2.5) is equivalent to

$$\begin{aligned} &\theta^2\alpha^2(x-kt)\psi^2(x-kt) + (1-\theta)^2\psi_x^2 - 2\theta(1+\theta)\alpha(x-kt)\psi(x-kt)\psi_x \\ &+ 4\varepsilon_1\theta\alpha(x-kt)\psi(x-kt)\psi_x \end{aligned}$$

$$\begin{aligned}
&= 2\theta(1 - \theta)\alpha(x - kt)\psi(x - kt)\psi_x - 2\theta(1 + \theta)\alpha(x - kt)\psi(x - kt)\psi_x \\
&+ 4\varepsilon_1\theta\alpha(x - kt)\psi(x - kt)\psi_x \\
&= (-4\theta + 4\varepsilon_1)\theta\alpha(x - kt)\psi(x - kt)\psi_x \leq 0.
\end{aligned} \tag{2.11}$$

Example 2. We may choose suitable functions $\alpha_-(x - kt)$ and $\alpha_+(x - kt)$ to obtain the functions $\psi(x - kt)$ satisfying

$$\begin{aligned}
&\theta^2\alpha_-^2(x - kt)\psi^2(x - kt) + (1 - \theta)^2\psi_x^2 - 2\theta(1 + \theta)\alpha_-(x - kt)\psi(x - kt)\psi_x \\
&+ 4\varepsilon_1\theta\alpha_-(x - kt)\psi(x - kt)\psi_x \leq 0
\end{aligned} \tag{2.12}$$

and

$$4(-k + (1 + \varepsilon_1)\psi(x - kt))\psi_x(x - kt) - \alpha_+(x - kt)\psi^2(x - kt) \geq 0 \tag{2.13}$$

or $\chi(x - kt)$ satisfying

$$\begin{aligned}
&\theta^2\alpha_-^2(x - kt)\chi^2(x - kt) + (1 - \theta)^2\chi_x^2 + 2\theta(1 + \theta)\alpha_-(x - kt)\chi(x - kt)\chi_x \\
&- 4\varepsilon_1\theta\alpha_-(x - kt)\chi(x - kt)\chi_x \leq 0
\end{aligned} \tag{2.14}$$

and

$$4(-k + (1 + \varepsilon_1)\chi(x - kt))\chi_x(x - kt) - \alpha_+(x - kt)\chi^2(x - kt) \geq 0 \tag{2.15}$$

for a small $\varepsilon_1 > 0$.

For instance, if we let $\alpha_-(x), \psi(x)$ satisfy (2.9) or

$$\psi_x(x - kt) = -\frac{\beta}{2}\alpha_-(x - kt)\psi(x - kt), \tag{2.16}$$

then (2.12) is true. Clearly (2.13) is also true if

$$4(-k + (1 + \varepsilon_1)M)\psi_x(x - kt) - M\alpha_+(x - kt)\psi(x - kt) \geq 0 \tag{2.17}$$

or

$$\psi_x(x - kt) \geq -l_0\alpha_+(x - kt)\psi(x - kt) \tag{2.18}$$

for a suitable positive constant l_0 . If we choose $\alpha_-(x - kt)$ and $\alpha_+(x - kt)$ such that

$$-\frac{\beta}{2}\alpha_-(x - kt) \geq -l_0\alpha_+(x - kt), \tag{2.19}$$

then $\psi(x - kt)$ satisfies (2.13).

The proofs of Theorems 1-3 are given in the following several sections.

3. Proof of Theorem 1

By using the transformation $\Upsilon = \psi(x, t) + \pi$, for a suitable function $\psi(x, t)$ in (1.10), we have

$$\begin{aligned} & \pi_t + \psi_t + \left(u - \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}\right)(\pi_x + \psi_x) \\ &= \varepsilon \pi_{xx} + \varepsilon \psi_{xx} + \frac{2\varepsilon}{\nu} \nu_x \pi_x + \frac{2\varepsilon}{\nu} \nu_x \psi_x - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') \nu_x^2 \\ & - \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} (\psi(x, t) + \pi - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu) + \zeta(x, t) u |u|, \end{aligned} \tag{3.1}$$

which is

$$\begin{aligned} & \pi_t + \psi_t + \left(u - \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}\right) \pi_x - \psi_x (\psi(x, t) + \pi - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu) - \psi_x \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \\ &= \varepsilon \pi_{xx} - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') [\nu_x^2 - \frac{4\nu \sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \nu_x \psi_x + (\frac{2\nu \sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x)^2] \\ & + \varepsilon \psi_{xx} + \frac{2\varepsilon}{\nu} \nu_x \pi_x + \frac{2\varepsilon \sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x^2 - \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \psi(x, t) \\ & - \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \pi + \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu + \zeta(x, t) u |u| \end{aligned} \tag{3.2}$$

or

$$\begin{aligned} & \pi_t + \psi_t + a(x, t) \pi_x + b(x, t) \pi + \left[-\frac{2\varepsilon \sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x^2 - \varepsilon \psi_{xx} - \varepsilon_1 \psi(x, t) \psi_x\right] \\ & + \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \psi_x - (1 - \varepsilon_1) \psi(x, t) \psi_x - \zeta(x, t) u |u| \\ & + [\alpha(x - kt) \psi(x, t) - \psi_x] (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \\ & - \alpha(x - kt) (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \leq \varepsilon \pi_{xx}, \end{aligned} \tag{3.3}$$

for a suitable small constant $\varepsilon_1 > 0$, where $a(x, t) = u - \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \frac{2\varepsilon}{\nu} \nu_x$ and $b(x, t) = -\psi_x + \alpha(x - kt) (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu}$.

Similarly, if we make the transformation $\Delta = \psi(x, t) + s$ in (1.9), we obtain

$$\begin{aligned} & s_t + \psi_t + \left(u + \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}\right) (s_x + \psi_x) \\ &= \varepsilon s_{xx} + \varepsilon \psi_{xx} + \frac{2\varepsilon}{\nu} \nu_x s_x + \frac{2\varepsilon}{\nu} \nu_x \psi_x - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') \nu_x^2 \\ & + \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} u - \zeta(x, t) u |u|, \end{aligned} \tag{3.4}$$

which is

$$\begin{aligned}
& s_t + \psi_t + \left(u + \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}\right) s_x + \psi_x (\psi(x, t) + s - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu) + \psi_x \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} \\
& = \varepsilon s_{xx} - \frac{\varepsilon}{2\nu^2 \sqrt{\sigma'(\nu)}} (2\sigma' + \nu\sigma'') [\nu_x^2 + \frac{4\nu\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \nu_x \psi_x + (\frac{2\nu\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x)^2] \\
& + \varepsilon \psi_{xx} + \frac{2\varepsilon}{\nu} \nu_x s_x + \frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x^2 + \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} u - \zeta(x, t) u |u|
\end{aligned} \tag{3.5}$$

or

$$\begin{aligned}
& s_t + \psi_t + c(x, t) s_x + d(x, t) s + \left[-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x^2 - \varepsilon \psi_{xx} - \varepsilon_1 \psi(x, t) \psi_x\right] \\
& + \psi_x \left(\frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu\right) + (1 + \varepsilon_1) \psi(x, t) \psi_x \\
& - \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} u + \zeta(x, t) u |u| \leq \varepsilon s_{xx},
\end{aligned} \tag{3.6}$$

where $c(x, t) = u + \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \frac{2\varepsilon}{\nu} \nu_x$ and $d(x, t) = \psi_x$.

By using the maximum principle to the first equation in system (1.6), we may obtain the a-priori estimate $\nu \geq 2\xi$. Then letting $\varepsilon = o(\xi)$ and $\varepsilon = o(\varepsilon_1)$, we have the following estimates on the three terms of the left-hand side of (3.3) and (3.6)

$$-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''} \psi_x^2 - \varepsilon \psi_{xx} - \varepsilon_1 \psi(x, t) \psi_x > 0. \tag{3.7}$$

Now we rewrite (3.3) and (3.6) as follows:

$$\begin{aligned}
& \pi_t + \psi_t + a(x, t) \pi_x + b(x, t) \pi \\
& + \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \psi_x - (1 - \varepsilon_1) \psi(x, t) \psi_x + \frac{1}{2} \zeta(x, t) (\pi - s) |u| \\
& + [\alpha(x - kt) \psi(x, t) - \psi_x] (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \\
& - \alpha(x - kt) (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \leq \varepsilon \pi_{xx},
\end{aligned} \tag{3.8}$$

and

$$\begin{aligned}
& s_t + \psi_t + c(x, t) s_x + d(x, t) s \\
& + \psi_x \left(\frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu\right) + (1 + \varepsilon_1) \psi(x, t) \psi_x \\
& + \frac{1}{2} (\zeta(x, t) |u| - \alpha(x - kt) \frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)}) (s - \pi) \leq \varepsilon s_{xx}.
\end{aligned} \tag{3.9}$$

If we may choose a suitable bounded function $\psi(x, t)$ such that the following inequalities hold

$$\begin{aligned}
& \psi_t + \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \psi_x - (1 - \varepsilon_1) \psi(x, t) \psi_x \\
& + [\alpha(x - kt) \psi(x, t) - \psi_x] (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \\
& - \alpha(x - kt) (\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \geq 0
\end{aligned} \tag{3.10}$$

and

$$\psi_t + \psi_x \left(\frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \right) + (1 + \varepsilon_1)\psi(x, t)\psi_x \geq 0, \tag{3.11}$$

then we have from (3.8) and (3.9) that

$$\pi_t + a(x, t)\pi_x + b(x, t)\pi + \frac{1}{2}\zeta(x, t)|u|(\pi - s) \leq \varepsilon\pi_{xx} \tag{3.12}$$

and

$$s_t + c(x, t)s_x + d(x, t)s + \frac{1}{2}(\zeta(x, t)|u| - \alpha(x - kt)\frac{\nu - 2\xi}{\nu}\sqrt{\sigma'(\nu)})(s - \pi) \leq \varepsilon s_{xx}. \tag{3.13}$$

Before we check the possibility of (3.10) and (3.11), we apply the inequalities (3.12) and (3.13) to prove the following Lemma 4 about the a priori estimates of π and s :

Lemma 4. *If at the time $t = 0$, $\pi(x, 0) \leq 0$ and $s(x, 0) \leq 0$, then the maximum principle is true to the functions $\pi(x, t)$ and $s(x, t)$, namely, $\pi(x, t) \leq 0, s(x, t) \leq 0$ for all $t > 0$.*

Proof of Lemma 4. Make a transformation

$$\pi = e^{\beta t} \left(\bar{\pi} + \frac{N(x^2 + qLe^t)}{L^2} \right), \quad s = e^{\beta t} \left(\bar{s} + \frac{N(x^2 + qLe^t)}{L^2} \right), \tag{3.14}$$

where L, q, β are suitable positive constants and N is the upper bound of π, s on $R \times [0, T]$ (N can be obtained by the local existence). The functions $\bar{\pi}, \bar{s}$, as are easily seen, satisfy the equations

$$\left\{ \begin{aligned} &\bar{\pi}_t + a(x, t)\bar{\pi}_x - \varepsilon\bar{\pi}_{xx} + (\beta + b(x, t) + \frac{1}{2}\zeta(x, t)|u|)\bar{\pi} - \frac{1}{2}\zeta(x, t)|u|\bar{s} \\ &\leq -(qLe^t + 2xa(x, t) - 2\varepsilon)\frac{N}{L^2} - (\beta + b(x, t))\frac{N(x^2 + qLe^t)}{L^2}, \\ &\bar{s}_t + c(x, t)\bar{s}_x - \varepsilon\bar{s}_{xx} + (\beta + d(x, t) + \frac{1}{2}\zeta(x, t)|u| - \alpha(x - kt)\frac{\nu - 2\xi}{\nu}\sqrt{P'(\nu)})\bar{s} \\ &- (\frac{1}{2}\zeta(x, t)|u| - \alpha(x - kt)\frac{\nu - 2\xi}{\nu}\sqrt{P'(\nu)})\bar{\pi} \\ &\leq -(qLe^t + 2xc(x, t) - 2\varepsilon)\frac{N}{L^2} - (\beta + d(x, t))\frac{N(x^2 + qLe^t)}{L^2}, \end{aligned} \right. \tag{3.15}$$

resulting from (3.12) and (3.13). Moreover

$$\bar{\pi}(x, 0) = \pi(x, 0) - \frac{N(x^2 + qL)}{L^2} < 0, \quad \bar{s}(x, 0) = s(x, 0) - \frac{N(x^2 + qL)}{L^2} < 0, \tag{3.16}$$

$$\bar{\pi}(+L, t) < 0, \quad \bar{\pi}(-L, t) < 0, \quad \bar{s}(+L, t) < 0, \quad \bar{s}(-L, t) < 0. \tag{3.17}$$

From (3.15), (3.16) and (3.17), we have

$$\bar{\pi}(x, t) < 0, \quad \bar{s}(x, t) < 0, \quad \text{on} \quad (-L, L) \times (0, T). \tag{3.18}$$

If (3.18) is violated at a point $(x, t) \in (-L, L) \times (0, T)$, let \bar{t} be the least upper bound of values of t at which $\bar{\pi} < 0$ (or $\bar{s} < 0$); then by the continuity we see that $\bar{\pi} = 0, \bar{s} \leq 0$ at some points $(\bar{x}, \bar{t}) \in (-L, L) \times (0, T)$. So

$$\bar{\pi}_t \geq 0, \quad \bar{\pi}_x = 0, \quad -\varepsilon \bar{\pi}_{xx} \geq 0, \quad \text{at} \quad (\bar{x}, \bar{t}). \quad (3.19)$$

If we choose sufficiently large constants q, β (which may depend on the bound of the local existence) such that

$$qL + 2xa(x, t) - 2\varepsilon > 0, \quad \beta + b(x, t) > 0 \quad \text{on} \quad (-L, L) \times (0, T). \quad (3.20)$$

(3.19) and (3.20) give a conclusion contradicting the first inequality in (3.15). So (3.18) is proved. Therefore, for any point $(x_0, t_0) \in (-L, L) \times (0, T)$,

$$\pi(x_0, t_0) < \left(\frac{N(x_0^2 + qLe_0^t)}{L^2}\right)e^{\beta t_0}, \quad s(x_0, t_0) < \left(\frac{N(x_0^2 + qLe_0^t)}{L^2}\right)e^{\beta t_0}, \quad (3.21)$$

which gives the desired estimates $\pi \leq 0, s \leq 0$ if we let L go to infinity. So Lemma 4 is proved.

From $v \leq 0, s \leq 0$, we may immediately obtain the estimates $\Delta \leq \psi(x, t)$ and $\Upsilon \leq \psi(x, t)$ given in Part (I) of Theorem 1, if we may choose $\psi(x, t)$ such that (3.10) and (3.11) are true.

Lemma 5. Let $\sigma(\nu) = \frac{1}{\gamma}\nu^\gamma, 1 < \gamma \leq 3, c = 0$. For a given function $\psi(x) \in \mathcal{B}_d^2(R)$, if $\psi(x) \leq M < k$ and satisfies the inequality (2.5) in Theorem 1, then (3.10) and (3.11) are true if we choose $\psi(x, t) = \psi(x - kt)$.

Proof of Lemma 5. We first prove (3.11). Let $\psi(x, t) = \psi(x - kt)$, then $\psi_t = -k\psi_x$. When $\sigma(\nu) = \frac{1}{\gamma}\nu^\gamma, 1 < \gamma \leq 3$ and $c = 0$, we have

$$\frac{\nu - 2\xi}{\nu} \sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu = (\nu - 2\xi)\nu^{\frac{\gamma-3}{2}} - \int_{2\xi}^\nu s^{\frac{\gamma-3}{2}} ds - \int_0^{2\xi} s^{\frac{\gamma-3}{2}} ds \leq 0, \quad (3.22)$$

and

$$\psi_t + (1 + \varepsilon_1)\psi(x, t)\psi_x = (-k + (1 + \varepsilon_1)\psi(x - kt))\psi_x \geq 0. \quad (3.23)$$

Thus (3.11) is proved.

To prove (3.10), we let the left side of (3.10) be L , then

$$\begin{aligned} L &= \left[\int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu - \frac{2\xi}{\nu} \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \right] \psi_x \\ &+ \left(\frac{2\xi}{\nu} \psi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu - k\psi_x \right) - (1 - \varepsilon_1)\psi(x - kt)\psi_x \\ &+ [\alpha(x - kt)\psi(x - kt) - \psi_x](\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \\ &- \alpha(x - kt)(\nu - 2\xi) \frac{\sqrt{\sigma'(\nu)}}{\nu} \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \\ &= -\frac{1}{\theta}\alpha(x - kt)(\nu - 2\xi)^2\nu^{2\theta-2} + (\alpha(x - kt)\psi(x - kt) - \psi_x + \frac{\psi_x}{\theta})(\nu - 2\xi)\nu^{\theta-1} \\ &+ \left(\frac{2\xi}{\nu} \psi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu - k\psi_x \right) - (1 - \varepsilon_1)\psi(x - kt)\psi_x - \frac{2\xi}{\theta}\alpha(x - kt)(\nu - 2\xi)\nu^{2\theta-2}. \end{aligned} \quad (3.24)$$

Since

$$-\frac{2\xi}{\theta}\alpha(x)(\nu - 2\xi)\nu^{2\theta-2} \geq 0 \quad (3.25)$$

and

$$\left(\frac{2\xi}{\nu}\psi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu - k\psi_x\right) = \left(\frac{2\xi}{\theta}\nu^{\frac{\gamma-3}{2}} - k\right)\psi_x \geq \left(\frac{(2\xi)^\theta}{\theta} - k\right)\psi_x \geq 0, \tag{3.26}$$

we have

$$\begin{aligned} L &\geq -\frac{1}{\theta}\alpha(x-kt)(\nu-2\xi)^2\nu^{2\theta-2} - (1-\varepsilon_1)\psi(x-kt)\psi_x \\ &+ (\alpha(x-kt)\psi(x-kt) - \psi_x + \frac{\psi_x}{\theta})(\nu-2\xi)\nu^{\theta-1} \\ &= -\frac{1}{\theta}\alpha(x-kt)[(\nu-2\xi)^2\nu^{2\theta-2} + \frac{\theta\psi_x - \psi_x - \theta\alpha(x-kt)\psi(x-kt)}{\alpha(x-kt)}(\nu-2\xi)\nu^{\theta-1} \\ &+ (\frac{\theta\psi_x - \psi_x - \theta\alpha(x-kt)\psi(x-kt)}{2\alpha(x-kt)})^2] - (1-\varepsilon_1)\psi(x-kt)\psi_x \\ &+ \frac{(\theta\psi_x - \psi_x - \theta\alpha(x-kt)\psi(x-kt))^2}{4\theta\alpha(x-kt)}. \end{aligned} \tag{3.27}$$

Since $\alpha(x-kt) \leq 0$ and $\psi(x-kt) > 0$, then $L \geq 0$ if we let $\psi_x \leq 0$ satisfy

$$-(1-\varepsilon_1)\psi(x-kt)\psi_x + \frac{(\theta\psi_x - \psi_x - \theta\alpha(x-kt)\psi(x-kt))^2}{4\theta\alpha(x-kt)} \geq 0, \tag{3.28}$$

which is equivalent to

$$\begin{aligned} &(\theta\psi_x - \psi_x - \theta\alpha(x-kt)\psi(x-kt))^2 \\ &\leq 4\theta\alpha(x-kt)\psi(x-kt)\psi_x - 4\theta\varepsilon_1\alpha(x-kt)\psi(x-kt)\psi_x \end{aligned} \tag{3.29}$$

or

$$\begin{aligned} &(\theta-1)^2\psi_x^2 - 2\theta(\theta+1)\alpha(x-kt)\psi(x-kt)\psi_x \\ &+ 4\theta\varepsilon_1\alpha(x-kt)\psi(x-kt)\psi_x + \theta^2(\alpha(x-kt)\psi(x-kt))^2 \leq 0. \end{aligned} \tag{3.30}$$

(3.30) is true for a suitably small $\varepsilon_1 > 0$ since the inequality (2.5) given in Theorem 1. **Part (I) of Theorem 1 is proved.**

To prove Part (II), we let $\Upsilon = \chi(x, t) + \pi, \Delta = \chi(x, t) + s$ for a suitable function $\chi(x, t)$ in (1.10) and (1.9), we may repeat the process in the proof of (3.3) and (3.6) to obtain

$$\begin{aligned} &\pi_t + \chi_t + a(x, t)\pi_x + b(x, t)\pi + \left[-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma' + \nu\sigma''}\chi_x^2 - \varepsilon\chi_{xx} + \varepsilon_1\chi(x, t)\chi_x\right] \\ &+ \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu\chi_x - (1 + \varepsilon_1)\chi(x, t)\chi_x - \zeta(x, t)u|u| \\ &+ [\alpha(x-kt)\chi(x, t) - \chi_x](\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \\ &- \alpha(x-kt)(\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \leq \varepsilon\pi_{xx}, \end{aligned} \tag{3.31}$$

and

$$\begin{aligned}
& s_t + \chi_t + c(x, t)s_x + d(x, t)s + \left[-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma'+\nu\sigma''}\chi_x^2 - \varepsilon\chi_{xx} + \varepsilon_1\chi(x, t)\chi_x\right] \\
& + \chi_x\left(\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\right) + (1 - \varepsilon_1)\chi(x, t)\chi_x \\
& - \alpha(x - kt)\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)}u + \zeta(x, t)u|u| \leq \varepsilon s_{xx},
\end{aligned} \tag{3.32}$$

where $\varepsilon_1 > 0$ is a suitable small constant.

With the help of Lemma 4, we only need to choose a suitable function $\chi(x, t) \in \mathcal{C}_d^2(\mathbb{R})$ and a constant c so that the following inequalities (which are similar with (3.10) and (3.11))

$$\begin{aligned}
& \chi_t + \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\chi_x - (1 + \varepsilon_1)\chi(x, t)\chi_x \\
& + [\alpha(x - kt)\chi(x, t) - \chi_x](\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \\
& - \alpha(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu \geq 0
\end{aligned} \tag{3.33}$$

and

$$\chi_t + \chi_x\left(\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\right) + (1 - \varepsilon_1)\chi(x, t)\chi_x \geq 0 \tag{3.34}$$

are correct.

The proof of the inequality (3.34) is simple because when $\gamma > 3$ and $k < 0$ satisfies the condition in (II) of Theorem 1, we may choose $\chi(x, t) = \chi(x - kt)$, $c = 0$ so that

$$\begin{aligned}
& \chi_t + \chi_x\left(\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\right) + (1 - \varepsilon_1)\chi(x, t)\chi_x \\
& = (-k + (1 - \varepsilon_1)\chi(x - kt) + (\nu - 2\xi)\nu^{\frac{\gamma-3}{2}} - \int_{2\xi}^\nu s^{\frac{\gamma-3}{2}}ds - \int_0^{2\xi} s^{\frac{\gamma-3}{2}}ds)\chi_x \\
& \geq (-k + (1 - \varepsilon_1)\chi(x - kt) - \frac{1}{\theta}(2\xi)^\theta)\chi_x \geq 0.
\end{aligned} \tag{3.35}$$

To prove (3.33), we let the left side of (3.33) be L_1 , then

$$\begin{aligned}
L_1 & = \left[\int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu - \frac{2\xi}{\nu} \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\right]\chi_x \\
& + \frac{2\xi}{\nu}\chi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu - k\chi_x - (1 + \varepsilon_1)\chi(x - kt)\chi_x \\
& + [\alpha(x - kt)\chi(x - kt) - \chi_x](\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} - \alpha(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu \\
& = -\frac{1}{\theta}\alpha(x - kt)(\nu - 2\xi)^2\nu^{2\theta-2} + (\alpha(x - kt)\chi(x - kt) - \chi_x + \frac{\chi_x}{\theta})(\nu - 2\xi)\nu^{\theta-1} \\
& + \frac{2\xi}{\nu}\chi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu - k\chi_x - (1 + \varepsilon_1)\chi(x - kt)\chi_x - \frac{2\xi}{\theta}\alpha(x - kt)(\nu - 2\xi)\nu^{2\theta-2}.
\end{aligned} \tag{3.36}$$

Since

$$\frac{2\xi}{\nu}\chi_x \int_0^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu \geq 0, \quad -\frac{2\xi}{\theta}\alpha(x)(\nu - 2\xi)\nu^{2\theta-2} \geq 0, \tag{3.37}$$

we have

$$\begin{aligned}
 L_1 &\geq -\frac{1}{\theta}\alpha(x-kt)(\nu-2\xi)^2\nu^{2\theta-2} + (\alpha(x-kt)\chi(x-kt) - \chi_x + \frac{\chi_x}{\theta})(\nu-2\xi)\nu^{\theta-1} \\
 &\quad -k\chi_x - (1 + \varepsilon_1)\chi(x-kt)\chi_x \\
 &= -\frac{1}{\theta}\alpha(x-kt)[(\nu-2\xi)^2\nu^{2\theta-2} + \frac{\theta\chi_x - \chi_x - \theta\alpha(x-kt)\chi(x-kt)}{\alpha(x-kt)}(\nu-2\xi)\nu^{\theta-1} \\
 &\quad + (\frac{\theta\chi_x - \chi_x - \theta\alpha\psi}{2\alpha})^2] - k\chi_x - (1 + \varepsilon_1)\chi\chi_x + \frac{(\theta\chi_x - \chi_x - \theta\alpha\psi)^2}{4\theta\alpha}.
 \end{aligned} \tag{3.38}$$

Since $\alpha(x-kt) \leq 0$ and $\chi(x-kt) > 0$, then $L_1 \geq 0$ if we let $\chi_x \geq 0$ satisfy

$$-k\chi_x - (1 + \varepsilon_1)\chi\chi_x + \frac{(\theta\chi_x - \chi_x - \theta\alpha\chi)^2}{4\theta\alpha} \geq 0, \tag{3.39}$$

which is equivalent to

$$(\theta-1)^2\chi_x^2 + \theta^2(\alpha\chi)^2 - 2\theta(\theta-1)\alpha\chi\chi_x - (4\theta k + 4\theta(1 + \varepsilon_1)\chi)\alpha\chi_x \leq 0 \tag{3.40}$$

or

$$\begin{aligned}
 &(\theta-1)^2\chi_x^2 + \theta^2(\alpha\chi)^2 + 2\theta(\theta+1)\alpha\chi\chi_x - 4\theta\varepsilon_1\alpha\chi\chi_x \\
 &- 4\theta\alpha\chi(k + (1 + \theta)\chi) \leq 0.
 \end{aligned} \tag{3.41}$$

Since $k + (1 + \theta)\chi \leq k + \frac{\gamma+1}{2}M < 0$ and the inequality (2.6) in (II) of Theorem 1, (3.41) is correct and so **Part (II) of Theorem 1 is proved.**

4. Proof of Theorem 2

When $\alpha(x-kt) = \alpha_-(x-kt) + \alpha_+(x-kt)$, where $\alpha_-(x-kt) \leq 0, \alpha_+(x-kt) \geq 0$, we may rewrite (3.3) as

$$\begin{aligned}
 &\pi_t + \psi_t + a(x,t)\pi_x + b(x,t)\pi + [-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma'+\nu\sigma''}\psi_x^2 - \varepsilon\psi_{xx} - \varepsilon_1\psi(x,t)\psi_x] \\
 &+ \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu\psi_x - (1 - \varepsilon_1)\psi(x,t)\psi_x - (\zeta(x,t)|u| + \alpha_+(x-kt)(\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu})u \\
 &+ [\alpha_-(x-kt)\psi(x,t) - \psi_x](\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \\
 &- \alpha_-(x-kt)(\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu \leq \varepsilon\pi_{xx}.
 \end{aligned} \tag{4.1}$$

In a similar way to obtain (3.6), we may obtain the following inequality

$$\begin{aligned}
 &s_t + \psi_t + c_1(x,t)s_x + d_1(x,t)s + [-\frac{2\varepsilon\sqrt{\sigma'(\nu)}}{2\sigma'+\nu\sigma''}\psi_x^2 - \varepsilon\psi_{xx} - \varepsilon_1\psi(x,t)\psi_x] \\
 &+ \psi_x(\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu) + (1 + \varepsilon_1)\psi(x,t)\psi_x \\
 &- \alpha_+(x-kt)(\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu}\psi(x,t) + \alpha_+(x-kt)(\nu-2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu}d\nu \\
 &+ (-\alpha_-(x-kt)\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} + \zeta(x,t)|u|)u \leq \varepsilon s_{xx},
 \end{aligned} \tag{4.2}$$

where $c_1(x, t) = c(x, t) = u + \frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \frac{2\xi}{\nu}\nu_x$ and $d_1(x, t) = \psi_x - \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu}$.

Therefore, if we may choose a suitable bounded function $\psi(x, t)$ such that the following inequalities hold

$$\begin{aligned} & \psi_t + \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \psi_x - (1 - \varepsilon_1)\psi(x, t)\psi_x \\ & + [\alpha_-(x - kt)\psi(x, t) - \psi_x](\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \\ & - \alpha_-(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \geq 0 \end{aligned} \quad (4.3)$$

and

$$\begin{aligned} & \psi_t + \psi_x \left(\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \right) + (1 + \varepsilon_1)\psi(x, t)\psi_x \\ & - \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu}\psi(x, t) \\ & + \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \geq 0, \end{aligned} \quad (4.4)$$

then we have from (4.1) and (4.2) that

$$\begin{aligned} & \pi_t + a(x, t)\pi_x + b(x, t)\pi \\ & + \frac{1}{2}(\zeta(x, t)|u| + \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu})(\pi - s) \leq \varepsilon\pi_{xx} \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} & s_t + c_1(x, t)s_x + d_1(x, t)s \\ & + \frac{1}{2}(\zeta(x, t)|u| - \alpha_-(x - kt)\frac{\nu-2\xi}{\nu}\sqrt{\sigma'(\nu)})(s - \pi) \leq \varepsilon s_{xx}. \end{aligned} \quad (4.6)$$

If we let $\psi(x, t) = \psi(x - kt)$, using the inequality (2.12), we may prove (4.3) in a similar way like the proof of (3.10).

Under the conditions in (III) of Theorem 2, (4.4) is true because

$$\psi_x \left(\frac{\nu - 2\xi}{\nu}\sqrt{\sigma'(\nu)} - \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \right) \geq 0 \quad (4.7)$$

and

$$\begin{aligned} & \psi_t + (1 + \varepsilon_1)\psi(x, t)\psi_x \\ & - \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu}\psi(x, t) \\ & + \alpha_+(x - kt)(\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu} \int_c^\nu \frac{\sqrt{\sigma'(\nu)}}{\nu} d\nu \geq (-k + (1 + \varepsilon_1)\psi(x - kt))\psi_x \\ & - \alpha_+(x - kt)\psi(x - kt)f(\nu) + \alpha_+(x - kt)f^2(\nu) \\ & = (-k + (1 + \varepsilon_1)\psi(x - kt))\psi_x - \frac{1}{4}\alpha_+(x - kt)\psi^2(x - kt) \end{aligned}$$

$$\begin{aligned}
 &7 + \alpha_+(f(\nu) - \frac{1}{2}\psi(x - kt))^2 \\
 &\geq (-k + (1 + \varepsilon_1)\psi(x - kt))\psi_x - \frac{1}{4}\alpha_+(x - kt)\psi^2(x - kt) \geq 0
 \end{aligned}
 \tag{4.8}$$

due to the condition (2.13), where $f(\nu) = (\nu - 2\xi)\frac{\sqrt{\sigma'(\nu)}}{\nu}$. Thus **Part (III) of Theorem 2 is proved**. Similarly we may prove Part (IV) of Theorem 2 and complete the proof of Theorem 2.

5. Proof of Theorem 3

From the upper estimates on the Riemann invariants given in Theorems 1-2, we can easily obtain the following estimates on $(\nu^{\xi,\varepsilon}, u^{\xi,\varepsilon})$,

$$2\xi \leq \nu^{\xi,\varepsilon}(x, t) \leq N(x, t), \quad |u^{\xi,\varepsilon}(x, t)| \leq N(x, t),
 \tag{5.1}$$

where $N(x, t)$ is a positive, bounded function, which depending on the bound of the initial data, but independent of ε, ξ .

Following the standard theory of semilinear parabolic systems, we can apply the contraction mapping principle to an integral representation of a solution to obtain the local existence result of the Cauchy problem (1.6)-(1.7). With the L^∞ estimate (5.1) of the local solution, we can extend the local time step by step to an arbitrary time T , since the step time depends only on the L^∞ norm.

As proved in [13], we know that the original system (1.1) and the approximated system (1.4) have the same entropy equation or the same entropies, and for any weak entropy-entropy flux pair $(\eta(\nu, u), q(\nu, u))$ of system (1.1)

$$\eta_t(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) + q_x(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t))
 \tag{5.2}$$

are compact in $H_{loc}^{-1}(R \times R^+)$, then the compactness framework given in [3,6] for $1 < \gamma < 3$ and in [7] for $\gamma \geq 3$ to ensure that there exists a subsequence of $(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t))$, which converges pointwisely to a pair of bounded functions $(\nu(x, t), u(x, t))$ as ξ, ε tend to zero, and the limit $(\nu(x, t), u(x, t))$ satisfies (2.3). Moreover, we multiply (1.6) by (η_ν, η_m) to obtain

$$\begin{aligned}
 &\eta_t(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) + q_x(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) + \xi q_{1x}(\nu^{\xi,\varepsilon}(x, t), u^{\xi,\varepsilon}(x, t)) \\
 &= \varepsilon \eta(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon})_{xx} - \varepsilon(\nu_x^{\xi,\varepsilon}, m_x^{\xi,\varepsilon}) \cdot \nabla^2 \eta(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon}) \cdot (\nu_x^{\xi,\varepsilon}, m_x^{\xi,\varepsilon})^T \\
 &+ \alpha(x - kt)u^{\xi,\varepsilon}m^{\xi,\varepsilon}\eta_\nu(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon}) \\
 &+ (\alpha(x - kt)u^{\xi,\varepsilon}m^{\xi,\varepsilon} - \zeta(x, t)m^{\xi,\varepsilon}|u^{\xi,\varepsilon}|)\eta_m(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon}) \\
 &\leq \varepsilon \eta(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon})_{xx} + \alpha(x - kt)u^{\xi,\varepsilon}m^{\xi,\varepsilon}\eta_\nu(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon}) \\
 &+ (\alpha(x - kt)u^{\xi,\varepsilon}m^{\xi,\varepsilon} - \zeta(x, t)m^{\xi,\varepsilon}|u^{\xi,\varepsilon}|)\eta_m(\nu^{\xi,\varepsilon}, m^{\xi,\varepsilon})
 \end{aligned}
 \tag{5.3}$$

for any weak convex entropy-entropy flux pair $(\eta(\nu, u), q(\nu, u))$ of system (1.1), where $q + \xi q_1$ is the entropy flux of system (1.4) corresponding to the entropy η . Thus the entropy inequality (2.4) is proved if we multiply a test function to (5.3) and let ε, ξ go to zero. Thus we obtain the proof of Theorem 3.

References

- [1] G.-Q. Chen, J. Glimm, Global solutions to the compressible Euler equations with geometric structure, *Commun. Math. Phys.* 180 (1996) 153–193.
- [2] S.-W. Chou, J.-M. Hong, B.-C. Huang, R. Quira, Global transonic solutions to combined Fanno Rayleigh flows through variable nozzles, *Math. Models Methods Appl. Sci.* 28 (2018) 1135–1169.
- [3] R.J. DiPerna, Convergence of the viscosity method for isentropic gas dynamics, *Commun. Math. Phys.* 91 (1983) 1–30.
- [4] C. Klingenberg, Y.-G. Lu, Existence of solutions to hyperbolic conservation laws with a source, *Commun. Math. Phys.* 187 (1997) 327–340.
- [5] D.I. Ketcheson, M. Parsani, R.J. Leveque, High-order wave propagation algorithms for hyperbolic systems, *SIAM J. Sci. Comput.* 35 (2013) A351–A377.
- [6] P.L. Lions, B. Perthame, P.E. Souganidis, Existence and stability of entropy solutions for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates, *Commun. Pure Appl. Math.* 49 (1996) 599–638.
- [7] P.L. Lions, B. Perthame, E. Tadmor, Kinetic formulation of the isentropic gas dynamics and p-system, *Commun. Math. Phys.* 163 (1994) 415–431.
- [8] T.P. Liu, Quasilinear hyperbolic systems, *Commun. Math. Phys.* 68 (1979) 141–172.
- [9] T.P. Liu, Nonlinear stability and instability of transonic gas flow through a nozzle, *Commun. Math. Phys.* 83 (1983) 243–260.
- [10] T.P. Liu, Resonance for a quasilinear hyperbolic equation, *J. Math. Phys.* 28 (1987) 2593–2602.
- [11] Y.-G. Lu, Global existence of solutions to system of polytropic gas dynamics with friction, *Nonlinear Anal., Real World Appl.* 39 (2018) 418–423.
- [12] Y.-G. Lu, Global existence of resonant isentropic gas dynamics, *Nonlinear Anal., Real World Appl.* 12 (2011) 2802–2810.
- [13] Y.-G. Lu, Some results on general system of isentropic gas dynamics, *Differ. Equ.* 43 (2007) 130–138.
- [14] N. Tsuge, Existence of global solutions for isentropic gas flow in a divergent nozzle with friction, *J. Math. Anal. Appl.* 426 (2015) 971–977.
- [15] N. Tsuge, Global L^∞ solutions of the compressible Euler equations with spherical symmetry, *J. Math. Kyoto Univ.* 46 (2006) 457–524.
- [16] N. Tsuge, Existence of global solutions for unsteady isentropic gas flow in a laval nozzle, *Arch. Ration. Mech. Anal.* 205 (2012) 151–193.
- [17] N. Tsuge, Isentropic gas flow for the compressible Euler equations in a nozzle, *Arch. Ration. Mech. Anal.* 209 (2013) 365–400.